

Claims

1. An RNA molecule derived from an alphavirus RNA genome and capable of efficient infection of animal host cells, which RNA molecule comprises the complete alphavirus RNA genome regions, which are essential to replication of the said alphavirus RNA, and further comprises an exogenous RNA sequence capable of expressing its function in said host cell, said exogenous RNA sequence being inserted into a region of the RNA molecule which is non-essential to replication thereof.

2. The RNA of claim 1, wherein the said alphavirus is Semliki Forest virus (SFV).

3. The RNA of claim 1 or 2, wherein the exogenous RNA sequence encodes a protein, a polypeptide or a peptide sequence defining an exogenous antigenic epitope or determinant.

4. The RNA of claim 3 wherein the exogenous RNA sequence encodes an epitope sequence of a structural protein of an immunodeficiency virus inclusive of the human immunodeficiency virus (HIV) types.

5. The RNA of any preceding claim, wherein the alphavirus derived RNA molecule regions comprise a 5' terminal portion, the coding region(s) for non structural proteins required for RNA replication, the subgenome promoter region and a 3' terminal portion of said viral RNA.

6. The RNA of claim 2, 3 or 5, wherein the exogenous RNA sequence encodes a foreign polypeptide or protein and is integrated into the SFV subgenomic 26S RNA substituting deleted parts thereof.

7. The RNA of claim 2, 3, 4 or 5, wherein the exogenous RNA sequence encodes a foreign viral epitopic peptide sequence and is located in a region of the RNA coding for structural alphavirus proteins enabling the exogenous RNA to be expressed as said viral epitope as part of the matured virus particle.

8. The RNA of claim 2, 3, 4 or 5, wherein the exogenous RNA sequence encodes a foreign viral epitopic peptide sequence inserted into the p62 spike precursor subunit encoding region of the SFV genome.

9. An RNA expression vector comprising the RNA of any preceding claim packaged into infectious particles comprising the RNA within the alphavirus nucleocapsid and surrounded by membrane with alphavirus spike proteins.

10. The vector of claim 9, wherein the RNA has a total size corresponding to the wild type alphavirus RNA genome or deviating therefrom to an extent compatible with package of the RNA into the infectious particles.

11. DNA transcription vector comprising a cDNA having one strand complementary to the RNA of any of claims 1 to 8.

12. A DNA expression vector comprising a full-length or partial cDNA complementary to alphavirus RNA or parts thereof and located immediately downstream of the SP6 RNA polymerase promoter and having a 5'ATGG or 5'GATGG or any other 5' terminus and a TTTCCA<sub>69</sub>ACTAGT or any other 3' terminus.

13. The vector of claim 12 having portions of the viral cDNA deleted, the deletions comprising the complete or part of the region(s) encoding the virus structural proteins, and further comprising an integrated polylinker region, which may correspond to BamHI-SmaI-XmaI, inserted at a location which enables an exogenous DNA fragment encoding a foreign polypeptide or protein to be inserted into the vector cDNA for subsequent expression in an animal host cell.

14. The vector of claim 12 or 13 wherein the alphavirus is SFV.

15. The vector of claim 12 or 14 comprising full-length cDNA and further comprising an exogenous DNA fragment encoding a foreign epitopic peptide sequence or antigenic determinant inserted into a region of the viral structural proteins.

16. The vector of claim 15 wherein the exogenous DNA fragment is inserted into the p62 spike precursor subunit encoding region of the SFV cDNA.

17. The vector of any preceding claim comprising an SFV derived cDNA which carries a conditionally lethal SFV mutation in the region encoding the p62 cleavage site, a cellularly uncleavable but extracellularly cleavable form of p62 being expressed.

18. The vector of claim 13 comprising SFV-derived cDNA, the vector being pSFV1, pSFV2 or pSFV3 having a structure as shown in Fig. 8.

19. An RNA transcript derived from transcription of the DNA-vector of any of claims 12-18 carrying an exogenous DNA fragment.

20. A method to produce the vector of claim 9 or 10

wherein the alphavirus derived RNA lacks part of or the complete region(s) encoding the structural viral proteins, the method comprising cotransfection of animal host cells with the RNA transcript of claim 19, wherein the alphavirus RNA lacks part(s) of or the complete region(s) encoding the viral structural proteins, with helper RNA transcribed in vitro from a helper DNA vector and culturing the host cells.

21. The method of claim 20 wherein the cotransfection is produced by electroporation of the host cells.

22. Helper vector for use in the method according to claim 20 or 21, said vector being comprised of the DNA vector of claim 12 wherein the regions encoding non structural virus proteins are almost completely deleted, including sequences encoding RNA signals for packaging of RNA into nucleocapsid particles, but the 5' and 3' signals needed for RNA replication and the region encoding the promoter for the structural sub-genome are in addition to those encoding the structural region preserved.

23. Helper vector of claim 22 wherein the cDNA has its origin from SFV and the deletion extends from the AccI (308) to the AccI (6399) restriction endonuclease site of the full-length cDNA vector of claim 12.

24. Helper vector of claim 22 and 23 where the structural region contains the mutation described in claim 17, or another conditionally lethal mutation.

25. The method of claim 20 wherein cells transformed to produce helper RNA according to claims 20, 22 or 23 are transfected with RNA transcript of claim 19.

26. A host cell of animal origin transformed with the RNA of any of claims 1-8, the DNA transcription vector of claims 11 or the DNA vector of any of claims 12-18 carrying an exogenous DNA fragment.

27. The host cell of claim 26 wherein the cell is an avian, a mammalian, a reptilian, an amphibian, an insecticidal or a fish cell.

28. The host cell of claim 27 which is the hamster BHK cell.

29. A method to produce the transformed host cell of claim 26, 27 or 28 comprising transfection of the cell with the RNA of any of claim 1-8, with the cDNA of claim 11 or of any of

claims 12-18 carrying an exogenous DNA fragment or infection of the cell with the infectious viral particles of claim 9 or 10.

30. The method of claim 29 wherein the transfection is produced by electroporation of the host cell.

31. A method for the production of a polypeptide or protein comprising infection of animal host cells with infectious particles according to claim 9 or 10, containing exogenous RNA encoding said polypeptide or protein and produced according to method of claim 20 or 21, culturing the said transformed cells to express the exogenous RNA and isolation and purification of the product formed by said expression.

32. A method for the production of a polypeptide or protein comprising in vitro transcription of the cDNA of the vector of any of claims 11-18 carrying an exogenous DNA fragment coding for the polypeptide or protein, transfection of animal host cells with the produced RNA transcript, transformed animal host cells being obtained harbouring the RNA transcript, culturing the said transformed cells to express the exogenous RNA and isolation and purification of the product formed by said expression.

33. The method of claim 32 wherein the vector cDNA is comprised of the cDNA of the vector of claim 17 carrying the exogenous DNA fragment.

34. An antigen consisting of a chimaeric alphavirus having an exogenous epitopic peptide sequence or antigenic determinant inserted into its structural proteins.

35. The antigen of claim 34 wherein the chimaeric alphavirus is derived from SFV.

36. The antigen of claim 34 or 35, wherein the exogenous epitopic peptide sequence is comprised of an epitopic peptide sequence derived from a structural protein of a virus belonging to the immunodeficiency virus class inclusive of the human immunodeficiency virus types.

37. Vaccine preparation comprising the antigen of claim 34, 35 or 36 as immunizing component.

38. Vaccine of claim 37 wherein the chimaeric alphavirus is attenuated by comprising the conditionally lethal SFV mutation of claim 17, an amber (stop codon) a temperature sensitive mutation or other mutation in its genome.

39. A method for the production of an antigen of claim

34, 35 or 36 comprising

a) in vitro transcription of the cDNA of the vector of any of claims 11-18 carrying an exogenous DNA fragment encoding the foreign epitopic peptide sequence or antigenic determinant and transfection of animal host cells with the produced RNA transcript, or

b) transfection of animal host cells with the said cDNA of the above step a),

culturing the transfected cells and recovering the chimaeric alphavirus antigen.

40. The method of claim 32, 33 or 39 wherein the transfection is produced by electroporation of the host cell.

41. A method for the production of an antigen in an organism by using in vivo infection with infectious particles according to claim 9 or 10 containing exogenous RNA encoding an exogenous epitopic peptide sequence or antigenic determinant, and produced according the claim 20 or 21.

-----

add B2

09901106-071001